



Huachenbio -- Your Professional Choice

苏州华辰生物科技有限公司

Suzhou Huachen Biotechnology Co.,Ltd

Tel: 400-965-9800

Website: www.huachenbiotech.com

Address: Unit A3-504-1, Creative Industry Park, No. 328 Xinghu Street,
Suzhou Industrial Park



WeChat
Official Account



WeChat Official
Subscription Account



WeChat ID
for Inquiries

- High expansion fold: >100,000-fold expansion by Day 28
- High cell quantity: approximately 200 billion cells by Day 21
- High purity: >98% CD3-CD56+ (High-Efficiency Version), >90% CD3-CD56+ (Enhanced Version), and >90% CD16+CD56+
- High cell viability: >90% viability
- Low cost: The production cost per NK cell preparation is 1/3 to 1/5 of the original cost
- Allogeneic use: CD3+ cells < 1%, meeting the requirement for allogeneic application
- Pure factor: feeder-free, GMP-manufactured, with DMF filing number




hyperClone® NK KIT

人NK细胞高效激活扩增试剂盒

hyperClone Human NK Activation/Expansion Cocktail

Brief Introduction to the Expansion Principle and Core Technology of hyperClone NK

HUACHEN hyperClone NK-kit integrates important signaling pathway factors essential for NK cell activation and expansion, as well as key cytokines that maintain and extend NK cell activity. Therefore, it can efficiently expand and activate high-purity NK cells, and serves as a powerful tool for large-scale preparation of NK cells with tumor-killing activity via the pure protein factor method.




hyperClone NK Cell Pure Protein Factor In Vitro Expansion Kit


It requires no feeder cells, enabling efficient expansion of NK cells with high purity to meet the application needs of autologous or allogeneic NK cells. With a defined chemical composition and compliance with GMP standards, it can support the regulatory filing of NK cell-based pharmaceuticals.

Sources of NK Cells

It is suitable for peripheral blood and umbilical cord blood, and compatible with both fresh blood samples and cryopreserved cells.




NK Cell Application and Operation Process





Data Presentation


Example 1: Growth and Expansion of Peripheral Blood-Derived NK (PB-NK) Cells




Typical Photos of Cell Proliferation in the First Three Days After Seeding



PBNK	DAY0	DAY6	DAY8	DAY9	DAY10	DAY12	DAY14	DAY16	DAY18	DAY20	DAY28
Calculate the total number of cells	8.98E+06	2.31E+08	6.90E+08	2.60E+09	4.47E+09	8.64E+09	1.84E+10	3.36E+10	6.67E+10	1.52E+11	8.99E+11
Expansion fold		26	77	290	498	962	2049	3742	7428	16927	100111
Volume after medium supplementation		450ml	900ml	1500ml	3L	6L	12L	24L	48L	96L	300L




Changes in Surface Marker Expression During PB-NK Cell Expansion




PBNK	7-AAD-Viability	CD45+	CD3-	CD3-CD56+	CD16+	CD56+CD16+	NKG2D	NKP44	NKP46	KIR	TRAIL
DAY6	98.86%	99.39%	97.99%	94.19%	46.51%	45.189%	45.86%	0.66%	51.29%	27.15%	1.370%
DAY8	98.17%	98.48%	99.68%	95.68%	64.75%	63.288%	90.26%	1.94%	54.18%	31.64%	0.347%
DAY10	93.69%	99.55%	98.53%	97.39%	87.82%	86.876%	91.13%	5.17%	68.99%	40.48%	1.917%
DAY14	99.59%	99.62%	99.96%	99.14%	96.84%	96.84%	90.88%	8.42%	72.95%	50.46%	0.200%
DAY16	99.78%	99.42%	99.88%	99.26%	96.96%	96.96%	90.53%	37.12%	77.26%	60.13%	2.265%
DAY21	99.16%	98.64%	99.99%	99.67%	98.24%	98.24%	93.15%	64.19%	79.78%	62.89%	1.275%

Example 2: Expansion Efficacy of CB-NK Cells




Detection of NK Cell Cytotoxicity


Tumor Killing Assay: Co-incubation of CBMCs with K562 Cells for 4 Hours After 14 Days of CBMC Culture



Tumor Killing Assay: Co-incubation of PBMCs with K562 Cells for 4 Hours After 14 Days of PBMC Culture




ADCC:RajiKilling ability Rituximab: 5 μ g/mL, co-incubation for 4 hours




Data Presentation


Cytokine and Cytotoxic Mediator Secretion: CD107a




Cytokine and Cytotoxic Mediator Secretion: TNF- α , IFN- γ



Cytokine and Cytotoxic Mediator Secretion: Granzyme B




Cytokine and Cytotoxic Mediator Secretion: Perforin




Allogeneic NK Cell Application—— Screening and Matching for Cord Blood NK Cell Bank Establishment

- CD16a 158V/V (The probability among East Asians is approximately 10%)
- HLA matching is not required
- Activating KIR, B-Type KIR, and Donor-Recipient KIR-HLA Mismatch



Allogeneic NK Cell Therapy — The Future of Cell Therapy Is Coming



The hyperClone NK Kit enables high-density, long-term culture and supports the acquisition of large quantities of allogeneic NK cell ready-to-use formulations with high purity and high activity.

Innovative Application Strategy Based on HUACHEN-NK Cells —Family NK Cell Healthcare



Family NK Cell Healthcare:

- ➊ A young member of the family donates a 50 mL blood sample;
- ➋ The laboratory performs expansion culture to establish 20-30 seed banks;
- ➌ Thawing one seed bank each time can support one NK cell healthcare session for a family member;
- ➍ Calculated based on two sessions per person per year, the seed banks can meet the family's needs for 15 years.
- ➎ Blood relatives who are young have cells with better activity, and these cells are more easily accepted (by the recipients' bodies).

Innovative Application Strategy Based on HUACHEN-NK Cells —NK Cell Healthcare for High-Net-Worth Individuals



NK Cell Healthcare for High-Net-Worth Individuals:

- ➊ First, adjust the body to an optimal state, then collect 50 mL of blood;
- ➋ Perform laboratory expansion and activation of the cells, followed by storage of 20-30 seed banks of cells;
- ➌ Retrieve one NK cell seed bank each time for two sessions annually, with two cell infusions per year;
- ➍ This setup can meet an individual's needs for 15 years.
- ➎ These are autologous cells, activated *in vitro* with high purity. In the future, they can support cell-based gene-modified therapy.

Top 10 Product Advantages

01	02	03	04
High Expansion Multiple With the pure factor kit, after 14-28 days of culture, the number of NK cells can be expanded by more than 100,000 times	High Purity, Double 90% Final Product: CD ⁻ CD56 ⁺ >95% CD16 ⁺ CD56 ⁺ >90%	High viability > 90%	Low CD3⁺ Ratio In the final product, CD3 ⁺ cells account for less than 5%, with most cases below 2%, which can meet the requirements for allogeneic infusion
05	06		
High Tumor-Killing Activity The killing rate against target cells reaches over 90% within 4 hours	10 Top 10 Product Advantages	Long In Vivo Survival Time Over 30 Days	
07	08	09	10
Simple Operation It can be completed without special equipment, is simple and easy to use, allowing researchers to easily obtain large quantities of high-purity NK cells after simple training	High Culture Success Rate Over 90% of samples can easily yield high-purity, qualified NK cells	Localization, Stable Supply Both cytokines and NK cell-specific culture media have achieved localization, enabling large-scale and stable supply	GMP - level, supporting the application for NK cell drugs The NK cell culture kits and dedicated culture media are produced in compliance with GMP standards and can meet the requirements for pharmaceutical registration

Guide for Selecting NK Cell Culture-Related Products

Product Name	Cat#	Main Parameters	Application Scenarios	Advantages
hyperClone® Human Fresh Blood NK Activation/Expansion Kit	hyperCloneNK-HCkit03	<ul style="list-style-type: none"> ① Pure cytokine kit, no feeder cells required; ② After 14-28 days of culture and expansion, the number of NK cells can be expanded by more than 100,000 times; ③ Purity can reach >98%, meeting the requirements for allogeneic infusion; ④ High CD16+CD56+ expression: when combined with antibody therapy, it can give full play to the antibody-dependent cell-mediated cytotoxicity (ADCC) effect of NK cells; ⑤ The kit adopts a 2L system; if large-scale expansion is needed, only the expansion reagent set needs to be supplemented, which can meet the demand for large-scale culture of NK cells at the 100-billion level; ⑥ Suitable for both peripheral blood and umbilical cord blood. 	1.Ultra-high purity NK cells, allogeneic NK cell infusion; 2.100-billion-level large-scale NK NK cell culture protocol using 50ml blood samples; 3.Meeting the requirements for NK cell drug IND filing.	Efficient and High-Purity Culture Restricted to Fresh Blood
hyperClone® Human Cryopreserved Mononuclear Cells NK Activation/Expansion Kit	hyperCloneNK-HCkit02	<ul style="list-style-type: none"> ① Specifically optimized for cryopreserved cells, compatible with cryopreserved PBMCs or cryopreserved CBMCs; ② For autologous NK cell use, the final purity of cryopreserved PBMC-derived NK cells is approximately 80-90%; ③ The final purity of cryopreserved CBMC-derived NK cells is over 90%; ④ Culturing in a 2L system yields 6-8 billion cells; ⑤ High cost-effectiveness. 	1.Preparing NK cells by resuscitating the client's cryopreserved peripheral blood PBMCs, suitable for autologous infusion; 2.Preparing NK cells by resuscitating cryopreserved CBMCs from umbilical cord blood banks (with the longest tested cryopreservation duration of 20 years for samples).	Efficient and cost-effective culture restricted to cryopreserved mononuclear cells
hyperClone® Human Fresh Blood NK Activation/Expansion Kit	hyperCloneNK-HCkit05	<ul style="list-style-type: none"> ① Specifically optimized for the preparation of NK cells from peripheral blood and umbilical cord blood in the field of general health; ② Pursuing high cost-effectiveness; ③ The final purity of peripheral blood-derived NK cells is >90%, for autologous use; ④ The final purity of umbilical cord blood-derived NK cells is >90%, suitable for allogeneic use. 	1.Fresh peripheral blood-derived NK cells for autologous use; 2.Preparation of fresh umbilical cord blood-derived NK cells;	1.Efficient and high-purity culture restricted to fresh blood; 2. Its purity is slightly lower than that of Kit 03, failing to meet the standard for allogeneic use of peripheral blood-derived NK cells, but its cost-effectiveness is significantly improved.
NK Cell Seed Bank Establishment, Resuscitation, and Recultivation Kit	Kit03+Kit09	<ul style="list-style-type: none"> ① Use the trial Kit 03 to culture 50ml of fresh blood sample for approximately Day 8 to Day 10, and establish a seed cell bank consisting of 20-30 aliquots, with each aliquot containing 1×10^8 cells (using StarSeed-02 as the cryopreservation solution); ② When needed, take one aliquot of seed cells, resuscitate and reculture them for about 10 days, and prepare one fresh NK cell preparation for clinical use. 	1.True NK cell storage with seed bank establishment; 2.Elimination of repeated blood collection; 3.Seed cells meeting the standards for allogeneic use; 4.Family-based NK cell health management program; 5.NK cell storage program for high-net-worth individuals.	One blood collection for multiple uses of fresh preparations
pureSep-NK NK Cell Purification Reagent	pureSep-01	<p>It enables simple and rapid purification of NK cells from whole blood, reduces T cell interference, requires no special operations or equipment, and adopts a non-magnetic bead method. It features high NK cell recovery rate and high cell viability.</p> <p>It uses a negative selection method, which does not affect the performance of NK cells.</p>	<p>It enables the purification of NK cells from umbilical cord blood or peripheral blood, eliminating individual differences in blood sample sources and ensuring that the purity of cultured NK cells reaches over 99%.</p>	<p>It is simple, rapid and efficient. Negative selection does not affect NK cell quality, eliminates individual differences in blood sample sources, and improves the success rate of NK cell culture.</p>
Cryopreservation Solution	StarSeed-01 StarSeed-02 HCCryo-GMP03 HCCryo-GMP04	<ul style="list-style-type: none"> ① Seed Cell Cryopreservation Solution: Features high cryopreservation and resuscitation efficiency, prevents organelle damage, and ensures a high success rate of cell culture after resuscitation; ② Final Preparation Cryopreservation Solution: A serum-free, animal-derived component-free, pharmaceutical-grade premixed ready-to-use cell cryopreservation solution, with a maximum cryopreservation density of 1×10^8 cells/mL 	<p>For PBMCs or CBMCs after Ficoll separation, StarSeed-01 Cryopreservation Solution shall be used;</p> <p>For cryopreservation of NK seed cells cultured until approximately Day 8, StarSeed-02 Cryopreservation Solution shall be used;</p> <p>For the final NK cell preparation, either GMP-03 Cryopreservation Solution (containing DMSO) or GMP-04 Cryopreservation Solution (DMSO-free) can be selected.</p>	<p>Select different cryopreservation solutions based on various cryopreservation purposes. These solutions are specially optimized and have clear functional divisions.</p>
novaNK-20 Chemically Defined Serum-Free Medium For Culture of Human Natural Killer cells	HC-NK20R HC-NK20F	<ul style="list-style-type: none"> ① Specifically developed for NK cell culture; ② Chemically defined; ③ Compared with similar products on the market, it enables faster cell proliferation, higher cell viability, and higher cell density; ④ The cell density in culture bags can reach $4-5 \times 10^6$ cells/mL. 	<p>It is suitable for culturing NK cells in culture bags and for large-scale culturing of NK cells in Wave bioreactors.</p>	<p>Higher cell density, faster proliferation, and higher viability.</p>

Huachen Bio: Core Technology Platforms



Ordering Information

Product Name	Cat#	Specs
hyperClone® Human Fresh Blood NK Activation/Expansion Kit	hyperCloneNK-HCkit03	1Kit, 2L
hyperClone® Human Cryopreserved Mononuclear Cells NK Activation/Expansion Kit	hyperCloneNK-HCkit02	1Kit, 2L
hyperClone® Human Fresh Blood NK Activation/Expansion Kit	hyperCloneNK-HCkit05	1Kit, 2L
hyperClone® NK Cells Seed Bank Resuscitation/Activation Culture Kit	hyperCloneNK-HCkit09	1Kit, 2L
hyperclone® NK Cells Amplification Reagent Kit	NKkit-20SE	1L
pureSep-NK NK Cell Purification Reagent	pureSep-01	1.5mL/tube
Cryopreservation Solution	StarSeed-01 StarSeed-02 HCCryo-GMP03 HCCryo-GMP04	100mL/bottle
novaNK-20 Chemically Defined Serum-Free Medium For Culture of Human Natural Killer cells	HC-NK20R HC-NK20F	1L/bottle

- Suitable for Primary Cell Isolation and Subculture
- Compatible with a variety of mesenchymal stem cells, such as umbilical cord, adipose-, bone marrow-, amniotic membrane-, hair follicle-, and dental pulp-derived mesenchymal stem cells
- Serum-free, free of any animal-derived components, antibiotic-free, stable in performance, and minimal batch-to-batch variation
- High cell expansion rate, with a single passage expansion fold of over 20x
- Cell yield per T175 flask: $>2 \times 10^7$ cells; Cell yield per 10-layer cell factory: $8-10 \times 10^8$ cells
- Cell diameter: 14-15 μ m, smaller than that of similar products on the market
- GMP level, prepared with water for injection (WFI), endotoxin <0.1 EU/ml
- Independent R&D and production system, stable supply, and high cost-effectiveness

- Serum-free, platelet lysate-free
- Animal-derived component-free
- Human-derived component-free, containing recombinant human serum albumin (rHSA)
- Chemically defined, with high batch-to-batch consistency
- High efficiency, supporting primary and subculture of MSCs
- Manufactured in compliance with GMP standards, supporting pharmaceutical registration
- Higher quality, with endotoxin <0.1 EU/ml



novastem-MSC®

间充质干细胞无血清培养基
Serum-Free Medium For Mesenchymal Stem Cell



- High expansion fold: >100,000-fold expansion by Day 28
- High cell quantity: approximately 200 billion cells by Day 21
- High purity: >98% CD3⁺CD56⁺ (High-Efficiency Version), >90% CD3⁺CD56⁺ (Enhanced Version), and >90% CD16⁺CD56⁺
- High cell viability: >90% viability
- Allogeneic use: CD3⁺ cells <1%, meeting the requirement for allogeneic application
- Low cost: The production cost per NK cell preparation is 1/3 to 1/5 of the original cost
- Pure factor: feeder-free, GMP-manufactured, with DMF filing number



hyperClone® NK KIT

人NK细胞高效扩增试剂盒
hyperClone Human NK Activation/Expansion Cocktail



StarMedium®

新一代间充质干细胞化学成分限定培养基
Chemically Defined Medium For Mesenchymal Stem cell



3D StarPore® Max

第二代Max微载体
3D StarPore gelatin dissolvable porous microcarrier for cell culture